

## 発表英語論文 (筆頭著者のみ)

1. Two-stage (posterior and anterior) surgical treatment using posterior spinal instrumentation for pyogenic and tuberculous spondylitis. Fukuta S, Miyamoto K, Masuda T, Hosoe H, Kodama H, Nishimoto H, Sakaeda H, Shimizu K. Spine (Phila Pa 1976). 2003 Aug 1;28(15):E302-8.
2. Unusual back pain caused by intervertebral disc degeneration associated with Schmorl node at Th11/12 in a young athlete, successfully treated by anterior interbody fusion: a case report. Fukuta S, Miyamoto K, Iwata A, Hosoe H, Iwata H, Shirahashi K, Shimizu K. Spine (Phila Pa 1976). 2009 Mar 1;34(5):E195-8.
3. High-grade(Grade III) Degenerative spondylolisthesis at L4/5 Treated Successfully by Transforaminal Interbody Fusion (TLIF): a case report. Fukuta S, Miyamoto K, Yoshida M, Kodama H, Kanamori Y, Hosoe H, Shimizu K. The Internet Journal of Orthopedic Surgery 2010, Vol16 Number 2
4. Kidney-type Intervertebral Spacers Should be Located Anteriorly in Cantilever Transforaminal Lumbar Interbody Fusion: Analyses of Risk Factors for Spacer Subsidence for a Minimum of 2 Years. Fukuta S, Miyamoto K, Hosoe H, Shimizu K. J Spinal Disord Tech. 2011 May;24(3):189-95.
5. Abundance of calpain and aggrecan-cleavage products of calpain in degenerated human intervertebral discs. Fukuta S, Miyamoto K, Suzuki K, Maehara H, Inoue T, Hara A, Kikuike K, Taguchi A, Shimizu K. Osteoarthritis Cartilage. 2011 Oct;19(10):1254-62.